

Plantlet Regeneration from Leaf Explants of *Gymnema Sylvestre* (Retz) R. Br. Ex Roemer & Schultes. An Important Medicinal Plant

Anitha Devi Uppu*

Head & Assistant Professor Department of Botany,
Govt. Degree & P.G. College for Women Karimnagar- 505001 (T.S). India.

Author for Correspondence: Uppu Anitha Devi

Abstract: The present study was undertaken to investigate the possibility for developing a one-step regeneration protocol in *Gymnema sylvestre* (Retz) R. Br. For this, the effect of Leaf explants was cultured on MS medium containing various concentrations of three cytokinins TDZ/BAP/Kn (0.5-3.0 mg/L). Maximum number of shoot bud proliferation was observed at (2.0mg/L). TDZ, compared to all other concentrations of Kn/BAP. As the concentration was increased above (2.0mg/L) the shoot bud induction was reduced gradually in TDZ/BAP/Kn. High frequency of shoots was induced at (2.0mg/L) TDZ. The *in vitro* regenerated shoots produced more number of roots on MS medium containing (1.0mg/L) IBA. Thus the plant developed *in vitro* using leaf cultures were established in pots containing garden soil outside under shade in wound temperature and light conditions. These plants flowered after 8 weeks following transfer to pots. The protocol established can be used for rapid multiplication of the specific producing true to type plants.

Keywords: *Gymnema sylvestre* (Retz) R. Br., multiplication, Leaf explants, TDZ, BAP, Kn and IBA.

Abbreviations: TDZ- Thidiazuron BAP- 6-Benzylaminopurin, Kn-Kinetin and IBA- Indole-3 butyric acid.

Date of Submission: 15-11-2017

Date of acceptance: 21-12-2017

I. Introduction

Gymnema sylvestre R.Br. which belongs to the family Asclepiadaceae, is a vulnerable species. It is a perennial slow growing woody climber of tropical and subtropical regions (Anonymous, 1997). It is a potent antidiabetic plant and used in folk, homeopathic and ayurvedic Systems of medicine (Mitra *et al.*, 1995). It is also used in treating of asthma, eye complaints, inflammations, family planning and snake bite (Anonymous, 1956; Selvanayagam *et al.*, 1995). In addition, it possesses antimicrobial, anti hypercholesterolemic (Bishayee and Chatterjee, 1994), hepatoprotective (Rana and Avadhoot, 1992) and sweet suppressing (Kurihara, 1992) activities. It also acts as feeding deterrents to caterpillar, *Prodenia eridania* (Granich *et al.*, 1974), prevent dental caries caused by *Streptococcus mutans* (Hiji Yasutake, 1990) and in skin cosmetics (Maeda *et al.*, 1996). There is a growing demand for leaves of *G. sylvestre* in the pharmaceutical trade due to its use as a remedy for diabetes and also as a tonic of the nerves and as a laxative (Shanmugasundaram *et al.*, 1983), as an anti-sweetner (Kurihara, 1992) and as an antihypercholesterolemic (Bishayee and Chatterjee, 1994). It also has stomatic, diuretic and cough suppressant property (Kapoor, 1990). Increasing awareness of the side effects of Western drugs have made general public turn towards the herbal medicine, thus the demands for medicinal plants have drastically increased. Due to over exploitation, this plant species has become threatened and is listed in International Union for Conservation of Nature (IUCN) red data book (Shailasree *et al.*, 2012).

Gymnema sylvestre is a slow growing, perennial woody climber (Shrivastava and Singh, 2011). Seeds lose viability in a short period of storage (Reddy *et al.*, 1998). Conventional propagation methods are hampered due to its poor seed viability, low rate of germination and poor rooting ability of vegetative cuttings (Komavali and Rao, 2000). The propagation of plant through seed results in less survivability under natural conditions (Anonymous, 1950). Therefore, to fulfill the increasing demand of this potent medicinal plant and population, *in vitro* culture and micropropagation could be an alternative method to aid in its conservation. Therefore, the propagation of this plant species by alternative methods is needed.

According to them, a maximum number of roots could be obtained on MS medium supplemented with NAA at 1.0 mg/L. (Karthic and Seshadri 2009) used young stem cuttings with an actively growing side branch as explants in their hydroponic experiment conducted with plastic tubes. MS medium containing 1/10 strength of MS salts supplemented with IBA at different concentrations (0.5, 1.0, 2.5 mg/L) and IBA (0.5 mg/L) resulted in the highest rooting (66%) and survival (96%). Based on the findings, they claimed that their protocol can

serve as an alternative to the existing *in vitro* and clonal multiplication protocols for *G. sylvestre*. (Subathra Devi and Srinivasan 2008) further stressed that *in vitro* propagation of *G. sylvestre* is not very different from that of *G. elegans* and may be applicable for other economically important woody cიმbers as well. Though direct regeneration studies have been conducted so far, this paper deals with the direct plant regeneration system with large number of shoots within a short period from leaf explants of *Gymnema sylvestre* induced by /TDZBAP/Kin.

II. Methodology

Plant material Seeds of *G. sylvestre* were collected from one single plant grown in the Forest range Officer Waddepally Hanamkonda. The seeds were washed in running tap water for 5 min, treated with 2-3 drops of Tween- 40 for 10 min and finally the seeds were washed thrice with sterile distilled water. Seeds were subsequently surface sterilized with 0.1% HgCl₂ for 5 min and washed thrice with sterile distilled water. The surface sterilized seeds were cultured on Murashige and Skoog (1962) medium supplemented with 2% sucrose and solidified with 0.8% agar (Himedia) (Ashok *et.al.*, 2002) pH of the medium was maintained at 5.8 samples were grown at a photoperiod of photo period under white fluorescent light of 40-60 mol m⁻² s⁻¹ intensity. For callus induction the explants viz, Leaf (0.8–1.0 cm²) explants from 8-week old seedlings were excised, these explants were inoculated to MS medium supplemented with various concentrations of (1.0- 3.0 mg/L) of cytokinins such as TDZ/BAP/Kn (Table-1) all the explant growth regulators were used as cytokines alone in culture media. All media were adjusted to pH 5.8 before addition of 0.8% agar agar and autoclaved at 121⁰C and 103 K pa for 20 minutes cultures in 25 x 150 mm cultures tubes.

Culture Media and Culture Conditions: The leaf explants approximately 0.5-10 cm from seedling of *G. sylvestre* were inoculated on to regeneration medium containing MS basal salts supplemented with different concentrations of TDZ/BAP/Kn. The pH of the media was adjusted to 5.8 either with 0.1 N HCL or 0.1N Na OH, solidified with 0.8% Difco-bacto agar and autoclaved at 121⁰C under 15 psi for 15-20 minutes. All the cultures were incubated at 25⁰C with 16h photo period under white fluorescent light of 40-60 mol m⁻² s⁻¹ intensity. The shoots proliferated from Leaf explants they were excised and cultured on a rooting medium consisting of MS medium supplemented with different (0.5-2.0 mg/L) IAA/IBA. The rooted plantlets were gently removed from the flasks and the roots were washed in tap water to remove traces of agar.

III. Acclimatization

Plants with roots were transferred during two weeks, after washing of the agar with distilled water and to pots with a mixture of soil rite (1:1). Potted plantlets were covered with transparent polythene membrane to ensure high humidity and watered every three days with half strength MS salts solution for two weeks in order to acclimatize plants to field conditions. After two weeks the acclimatized plants were transferred to pots containing normal garden soil and maintained in greenhouse under natural day length conditions.

IV. Results And Discussion

The results of the Leaf explants cultures on the development of multiple shoots and roots are shown in (Table-1). The Leaf explants of *G. sylvestre* cultured on different hormonal combinations showed varied results.

Table – 1: Influence of various concentrations of MS+ TDZ/BAP/Kin on induction of shoot buds proliferation from Leaf explants cultures of *G. sylvestre*.

Hormone concentration (mg/L)	% of frequency of plantlet production	Average No. of shoots / Explants ± (SE)*	Average length of shoots (cms) ± (SE)*
TDZ			
0.5	50	4.9 ± 0.42	1.0 ± 0.28
1.0	60	5.3 ± 0.07	2.4 ± 0.33
1.5	64	8.5 ± 0.62	3.4 ± 0.62
2.0	74	9.3 ± 0.43	3.6 ± 0.42
2.5	52	6.0 ± 0.25	2.3 ± 0.35
3.0	50	2.0 ± 0.25	2.0 ± 0.25
BAP			
0.5	48	3.2 ± 0.32	1.3 ± 0.38
1.0	53	4.2 ± 0.27	2.0 ± 0.23
1.5	58	4.5 ± 0.32	2.4 ± 0.42
2.0	60	8.3 ± 0.23	2.8 ± 0.12
2.5	50	4.2 ± 0.23	2.0 ± 0.25
3.0	42	2.0 ± 0.34	1.2 ± 0.35
Kin			
0.5	43	2.0 ± 0.52	1.2 ± 0.22

1.0	47	4.3 ± 0.37	1.8 ± 0.23
1.5	52	4.5 ± 0.42	2.0 ± 0.42
2.0	58	6.0 ± 0.23	2.2 ± 0.32
2.5	48	3.4 ± 0.65	1.8 ± 0.25
3.0	40	2.0 ± 0.45	1.1 ± 0.35

* Mean ± Standard Error

The Leaf explants became active with in weak after inoculation and new shoots became distinct by the seconds and third weak with leaves and internodes. The explants survival from leaf explants of nature plant of *G. sylvestre* varied with season. According to the present observations, the explants were collected from field grown plants thought out the year to determine the ideal season for culture established.

Table- 2 Influence of various concentrations of IAA/IBA on rooting of *in vitro* shoots of *G. sylvestre* after 8 weeks of incubation

Growth Hormones (mg/L)		Percentage of response	Average no of roots (S.E)*
IAA	IBA		
00	00	23	1.0 ± 0.12
0.5	-	60	2.3 ± 0.37
1.0	-	70	3.2 ± 0.38
2.0	-	70	5.6 ± 0.38
-	0.5	54	4.3 ± 0.36
-	1.0	73	8.3 ± 0.87
-	2.0	70	6.3 ± 0.36

* Mean ± Standard Error

Effect of TDZ

The leaf explants were excised from the surface sterilized, *in vitro* grown, 30-d old seedlings and cultured on MS medium augmented with TDZ (1.0–3.0 mg/L) for multiple shoot induction of all the different concentrations of TDZ tested, (2.0 mg/L) TDZ was found to be more effective in inducing (15.0 ± 0.3 shoots/explants). But at high concentration of TDZ (3.0 mg/L) considerably the number of shoot induction was found to be reduced. As the concentration of TDZ was increased up to 2.0mg/L the multiple shoots number was increased but as the concentration of TDZ (2.0mg/L) to (3.0 mg/L) TDZ resulted the number of shoots were reduced. (Fig-I a).

Effect of BAP

Leaf explants were cultured on MS medium amended with various concentrations of BAP (1.0-3.0 mg/L) as role growth regulators showed the direct organogenesis (Table -1) Maximum number of shoot bud proliferation (8.3 ± 0.23 shoots /explant) was found at (2.0 mg/L) BAP. At 1.0, and 2.0 mg/L BAP induced (3.2 ± 0.32 and 4.2 ± 0.27) shoots/explant with 48 and 53% of cultures responded. When the concentration was increased up to (3.0 mg/L) BAP gradually induction of multiple shoots were reduced. At (2.5 and 3.0 mg/L) BAP induced (4.2 ± 0.23 and 2.0 ± 0.34) shoots/explants with 50 and 42 percentage of cultures were responded. The number of shoot bud induction was found to be decreased as the concentration of BAP increased. At high concentration of BAP showed less number of shoots per explants. (Fig-I b).

Effect of Kn

Leaf explants were cultured on MS medium containing different concentrations of cytokinin Kn (1.0-3.0 mg/L) as role growth regulators showed the direct organogenesis/ shoot formation (Table -1) to find out the difference between TDZ, BAP and Kn in inducing the direct plant regeneration from Leaf explants in *G. sylvestre*. Maximum number of shoot bud proliferation (06.0 ± 0.23 shoots/ explant) was found at (2.0 mg/L) compared to all other concentrations of Kn at (1.0 and 2.0 mg/L) Induced (2.0 ± 0.52 and 4.3 ± 0.37 mg/L) shoots / explant with 438 and 47 percentage of cultures responded. Percentage of response was gradually increased up to (2.0 mg/L) Kn. At above (2.0 mg/L) concentration less number of shoots was recorded. At (2.5 and 3.0 mg/L) Kn induced (3.4 ± 0.65 and 2.0 ± 0.45) shoots/explants with 48 and 40 percentage of cultures were also recorded on MS + Kn. However high induction of ability was found in all the concentrations of TDZ, Kn compared to BAP.

In vitro rooting

Fully elongated healthy shoots were transferred on to full strength MS root induction medium (RIM) (Murashige and Skoog 1962) fortified with different concentration of IAA/IBA (0.5 – 2.0 mg/L). Profuse rhizogenesis was observed on (1.0 mg/L) IBA, compared to (0.5 -2.0 mg/L) IAA on MS medium containing (1.0 mg/L) IBA whereas 73% of plants produced roots with (8.3 ± 0.87) roots/ explant. (Fig-I -c).

V. Acclimatization

Rooted plantlets were removed from the culture medium and the roots were washed under running tap water to remove agar. Then the plantlets were transferred to polypots containing pre-soaked vermiculite and maintained inside a growth chamber set at 28° C and 70 – 80 % relative humidity. After three weeks they were transplanted to poly bags containing mixture of soil + sand + manure in 1: 1: 1 ratio and kept under shade house for a period of three weeks. The potted plantlets were irrigated with Hogland's solution every 3 days for a period of 3 weeks. (Fig-I d).




Fig-I Plantlet regeneration from Leaf explants of *Gymnema sylvestre* a), multiple shoot production on MS+(2.0mg/L) TDZ after six weeks (b) Plantlet proliferation on MS (2.0mg/L) BAP (c) root induction from leaf regenerate shoots on MS+(2.0mg/L) IBA (d) Hardening of plantlet.

VI. Discussion

We were successful in direct regenerating plants from leaf explants of *G. sylvestre* cultures on MS medium fortified with different concentrations of cytokinins i.e. TDZ/BAP/Kn (1.0-3.0 mg/L). Maximum number of shoot buds was induced at (2.0 mg/L) TDZ in comparison to Kn/BAP as role growth regulators. It was interesting find out that the shoots induction was enhanced in all the concentrations of cytokinins. However

the shoot bud proliferation was found to more on (2.0 mg/L) with TDZ compared to (2.0 mg/L) in with Kn/BAP but the concentration of TDZ induced highest number of plantlet regeneration among all hormonal combinations and concentrations were used in *G. sylvestre*.

The effect of TDZ was found to be highest the concentration of (2.0 mg/L) recording shoot buds/explants (9.3 ± 0.43). At lowest concentration of (0.5 mg/L) recording shoot buds/explants (4.9 ± 0.42) (Table-1). The average length of shoots and number of roots were found to be significantly higher in MS medium, supplemented with hormones. For Leaf explants *G. sylvestre*, the best growth regulator concentration for direct shoot induction in MS was observed in the presence of (2.0 mg/L) TDZ which gave the highest shoot elongation (2.8 ± 0.22). The presence of (0.1 mg/L) NAA and (2.0 mg/L) TDZ was best medium for shoot induction from nodal explants (Amarasinghe *et al.*, 2011). Which conform the results of the present study of Leaf explants. The presence of higher concentrations of BAP (2.0 mg/L) in the medium suppresses the shoot development (Komalavalli and Rao 2000) confirming the results of the present study, where the mean shoot length was drastically reduced in the presence of (1.5 mg/L) BAP. However, the growth regulator free MS (control) also showed high shoot elongation (14.2 ± 0.37) and the difference between those two treatments is not significant.

The results are summarized in (Table -2). Regenerated shoots were rooted on MS medium without supplementing any growth regulator (Reddy *et al.*, 1998). However, in contrast present study indicated that the growth regulator frees MS (Control) did not induce roots *in vitro*. ANOVA using Kruskal - Wallis Test indicated that all the treatments were significantly different on each other. High frequency of rooting (58%) was obtained in leaf explants derived shoots on MS medium supplemented with IBA (2.0 mg/L) (Komalavalli and Rao 2000). In the present study the best growth regulator concentration for *in vitro* root induction was observed in MS supplemented with (1.0 mg/L) IBA giving 73% success in root induction (Fig-1). This indicates that the decreased levels of basal salt concentrations may have affected the percentage rooting (Komalavalli and Rao 2000). Presence of lower levels of IBA (≤ 2.0 mg/l) did not induce roots in *in vitro* propagated shoots. Increased concentrations of higher than (1.0 mg/L) IBA not only retarded the root induction but also induced callus at the base of the shoot. Thus, it could be suggested that MS supplemented with (1.0 mg/L) IBA in MS would be the best medium for rooting of *in vitro* shoots of *G. sylvestre*.

The highest percentage of survival (73) was observed in the substrate with a high proportion of sand (soil: sand: 1: 2) (Table-2). (Reddy *et al.*, 1998) reported that for acclimation, plantlets were removed from the rooting medium 8 weeks after root initiation, and transferred to fresh tubes containing autoclaved tap water. After 8- 10 days, plantlets were subsequently transferred to plastic pots (9 × 9 cm) containing autoclaved soil rite covered with perforated polythene bags to maintain humidity; and were kept under culture room conditions for about 7 days. After three weeks, polythene bags were removed and pots were transferred to the garden, and placed under the shade till the new leaves appeared. Then they were planted under normal garden conditions. The transplantation success was about 75%. In the present study, substrate mixture with soil: sand (1 : 3) enhanced the surviving ability (Fig-1). An increasing portion of sand in the mixture enhances the surviving ability as smaller amount of water was retained in the substrate (Senarath *et al.* 2007). Plantlets are unable to compete with soil microbes and to cope with the harsh environmental conditions especially immediately after removal of the plantlets from *in vitro* conditions. After removal of the plantlets from the propagator, they were exposed to natural environment gradually. Yet, it was observed that especially during the rainy season, the survival ability of the plantlets was poor thus making it essential to keep the plantlets in a propagator for a long period if the rainy season persisted.

Acknowledgement

We thank to The University Grants Commission South Eastern Regional Office (SERO) Chirag-Ali-Lane Hyderabad- 500001, Hyderabad for providing the funds under the Minor Research Project (2015-2017) program to the Department of Botany Govt. Degree & P.G. College for Women Karimnagar. -505001, Telangana State India.

References

- [1]. Amarasinghe KIC, Silva TD and Weerahawa HLD (2011) *In vitro* propagation of *Gymnema sylvestre*. Proc. Ann. Acad. Ses. of Open University of Sri Lanka. pp. 245 - 248.
- [2]. Anonymous Medicinal Plants Categorized W/new IUCN Red List Criteria under the Biodiversity onservation Priorisation Project. In: 1997. CBSG, India News (pp. 10).
- [3]. Anonymous., 1956. The Wealth of India Raw Materials, Vol. IV (pp.276–277). Publication and Information Directorate. CSIR, NewDelhi.
- [4]. Bishayee A and Chatterjee M., 1994. Hypolipidaemic and antihypercholesterolemic effects of oral *Gymnema sylvestre* R. Br. Leaf extract in albino rats fed on a high fat diet. *Phytothera. Res.* 8: 118–120
- [5]. Granich S M., Halperin B P & Eisner T., 1974. Gymnemic acids: Secondary plant substances of dual defensive action. *J. Insec. Physiol.* 20: 435–439.
- [6]. Hiji Yasutake U S., 1990. Gymnemic acid for prevention of dental caries. In: *Chemical Abst.* Vol. 113, No. 46364b

- [7]. Kapoor LD., 1990. CRC Handbook of Ayurvedic medicinal plants: Boca Raton, Fl. pp: 200 - 201.
- [8]. Karthic R., Seshadri S., (2009.) Cost effective mass multiplication of *Gymnema sylvestre* in hydroponic system. Nature Precedings, hdl:10101/npre.2009.3770.1
- [9]. Komalavalli N & Rao MV., 2000. In vitro micropropagation of *Gymnema sylvestre* – A multipurpose medicinal plant. Plant Cell, Tissue and Organ Culture. 61: 97–105
- [10]. Kurihara Y., 1992. Characteristics of antisweet substances, sweet proteins and sweetness-inducing proteins. Crit. Rev. Food Sci. Nutr. 32: 231–252
- [11]. Maeda N and Fukuda M., 1996. Skin lightening cosmetics containing *Gymnema saponins* and other ingredients. In: Chemical Abst. Vol. 125, No. 123289h
- [12]. Mitra SK., Gopumadhavan S., Muralidhar TS., Anturlikar SD and Sujatha MB., 1995. Effect of D-400 herbomineral preparation on lipid profile glycated haemoglobin and glucose tolerance in streptozotocin induced diabetes in rats. Indian J. Exp. Biol. 33: 798–800.
- [13]. Murashige, T. and Skoog, F. A revised medium for rapid growth and bioassays with tobacco tissue culture. Physiol. Plant, 1962; 15: 473–497.
- [14]. Rana AC and Avadhoot Y., 1992. Experimental evaluation of hepatoprotective activity of *Gymnema sylvestre* and *Curcuma zedoaria*. Fitoterap. 63: 60–62.
- [15]. Reddy PS., Gopal GR and Sita GL., 1998. In vitro multiplication of *Gymnema sylvestre* R. Br. – An important medicinal plant. Curr. Sci. 75: 843–845
- [16]. Selvanayagam Z E., Gnanavendhan S G., Chandrasekharan P., Balakrishna K & Rao RB., 1995. Plants with antisnake venom activity-a review on pharmacological and clinical studies. Fitoterap. 65: 99–111.
- [17]. Senarath WTPSK, Chandrasena KGPH and Fernando KMEP (2007) In vitro mass propagation and greenhouse establishment of *Munronia pinnata* (Wall) Theob. Binkohomba. J. Nat. Sci. Found. 35(3): 181 - 189.
- [18]. Shailasree S., Ruma K., Kini KR., Niranjana SR and Prakash HS., 2012. Potential anti-inflammatory bioactives from medicinal plants of Western Ghats, India, J. Pharma. Com. 2(2): 12
- [19]. 12
- [20]. Shanmugasundaram KR., Panneerselvam C., Samudram P and Shanmugasundaram ER., 1983. Enzyme changes and glucose utilisation in diabetic rabbits: the effect of *Gymnema sylvestre*, R.Br. J. Ethnopharmacol. 7:205-234.
- [21]. Shrivastava R and Pratibha S., 2011. In vitro propagation of multipurpose medicinal plant *Gymnema sylvestre* R.Br. (Gudmar), Ind. J. Shodha anusandhan Samachar 1: 27 30.
- [22]. Subathra Devi C., Mohana A.V., In vitro propagation of *Gymnema sylvestre*. Asian Journal of Plant Sciences, 2008; 7: 660-665.

International Journal of Engineering Science Invention (IJESI) is UGC approved Journal with Sl. No. 3822, Journal no. 43302.

Anitha Devi Uppu*. “Plantlet Regeneration from Leaf Explants of *Gymnema Sylvestre* (Retz) R. Br. Ex Roemer & Schultes. An Important Medicinal Plant.” International Journal of Engineering Science Invention, vol. 06, no. 12, 2017, pp. 32-37.