

L.Minor used as a tool for Evaluation of Cu- Pollution in bio-monitoring program

Dr. HimansuBhusana Panigrahy¹, Tapan Panda², Sarala Prasanna Patnaik³

^{1,2}Gandhi Institute for Technology, Bhubaneswar

³Gandhi Engineering College, BBSR.

ABSTRACT: Macrophytes can be used as biomonitoring tools as they accumulate various heavy metals in their biomass. In this study, the impact of copper on the growth of the Lemna minor. Its removal, was also studied with 1 mg/L of Cu in a quarter of Lesaint solution maintained at 6.1pH. In order to verify the weed's tolerance to Cu, photosynthesis was measured at the maximal concentration which caused no effect on the plant growth. The results showed that copper inhibited Lemna growth at concentrations ≥ 0.3 mg/L. At 0.2 mg/L, the final biomass was approximately four times greater than the initial biomass. Analysis of metal concentration in water showed that Lemna minor was responsible for the removal of 26% of Cu from the solution. In the presence of Cu, respiration was reduced, while photosynthesis increased considerably. Net photosynthesis approximately increased three times compared to the control. Copper was responsible for 130-290% increase in the photosynthetic activities. These results suggested that Lemna minor could be a good tool for the evaluation of copper pollution in biomonitoring programs.

Key words: Aquatic pollution; Bio-indicator; Physiological modifications; Growth; Photosynthesis

I. Introduction

Copper (Cu) is an essential element for organisms and is involved in numerous physiological processes (Teisseire and Guy, 2000). However, it is toxic at higher concentrations by causing deleterious effects to human, animals and plants (Vinodhini and Narayanan, 2009). Excess of Cu may reach living organisms as a result of environmental pollution caused by anthropogenic activities (mining operations, manufacturing industries and agricultural technologies) which can modify the biogeochemical cycles of the metal.

Several studies demonstrated that many species of duckweed, a group of free-floating freshwater plants of the family Lemnaceae, are able to absorb and accumulate high amount of copper in their biomass producing an internal concentration several fold greater than the nutrient medium (Jain et al., 1989; Zayed et al., 1998; Miretzky et al., 2004; Ater et al., 2006). This accumulation has, in some ways, a relationship with the tolerance phenomena which is defined as the cell capability to protect plant tissues against injury caused by the metal (Sabreen and sugiyama, 2008). At metal concentration greater than the tolerated concentration, toxicity symptoms and physiological changes are induced.

Cupric ions are responsible for many alterations of plant cells and inhibition of enzymatic activities (Teisseire and Guy, 2000). They also cause significant changes in respiration, photosynthetic CO₂ fixation and photosynthetic pigments by increasing oxidation of chloroplast membranes (Prasad et al., 2001; Hattabet et al., 2009). These physiological modifications, evaluated by biotoxicity tests, can be used as an indicator of metal pollution and offer data in biomonitoring (Movahedian et al., 2005).

analysis was carried out with an atomic absorption spectrophotometer which had a detection limit of 10⁻² mg/L (Perkin Elmer).

The elimination percentage of Cu was calculated according to Khellaf and Zerdaoui (2009b):

$$\text{Elimination(\%)} = \frac{C_0 - C_f}{C_0} \times 100 \quad (1)$$

C₀-C_f Sedentary macrophytes as bioindicators have some advantages such as high tolerance to aquatic metal pollution, convenience for sampling, large individuals and easy to realize laboratory raise (Zhou et al., 2008). Duckweed has been widely used in toxicity tests of different chemicals and effluents and particularly, Lemna minor has often been selected to represent vascular aquatic plants in toxicity tests (Kanoun-Bouléet et al., 2009). This genus is an invasive plant wild-growing in European regions and other Mediterranean countries.

The main objective of the present study was to evaluate the effects of elevated Cu levels on Lemna minor growth and photosynthesis. The specific objectives were to: (a) determine the growth of plants in experimental Cu treatments ranging from 0.1 to 1.0 mg/L, (b) assess Cu removal percentage in the presence of

duckweed and (c) verify the plant tolerance to Cu by measuring net photosynthesis and respiration at the maximal concentration which causes no effect on the plant growth. The results were compared with those of other studies traditionally reported in literature.

II. Materials And Methods

Plant Toxicity Test And Metal Removal

The effect of copper on the growth of duckweed was assessed according to the test protocols derived from the standard draft guideline 221 of the Organization for Economic Cooperation and Development (OECD, 2002). The details of data analysis were the same as those described in our previous study (Khellaf and Zerdaoui, 2009a).

Water samples (1 mL) were regularly drawn in order to ascertain the Cu concentration removed from the solution. The metal concentration in which: C_0 and C_f are initial and remaining concentrations of metal in the medium (mg/L).

photosynthesis experiment

The essays were investigated using an infrared gas analyzer (IRGA) in a closed system with an airflow of 1.1×10^{-2} mL/min (the capacity of the circuit and the room of assimilation was of 11 litres). A 42-cm² frond area (previously exposed to 0.2 mg/L of Cu during 4 days) contained in a crystallising cup with a low volume of water was placed in an enclosure of which the atmosphere was renewed permanently (Garry, 1988). At this concentration, no morphological sign of toxicity was observed on Lemna fronds. Control treatment corresponded to the same essay without exposing plants to Cu. The infrared analyser used in differential mode allowed the direct measurement of the difference in CO₂ concentration between the entry and the exit. These measurements were taken with regular time intervals. The CO₂ concentration in the air (370 mg/L) was used for the calibration of the apparatus.

The plants were initially placed in the darkness (respiration 1); then successively exposed to three lamps (1st, 2nd and the 3rd photosynthesis) and replaced finally in the darkness (respiration 2). The duration of essays was 100 min corresponding to a time of 20 min for each phase (2 phases of respiration and 3 phases of photosynthesis). Photosynthetic active irradiations of the 1st, 2nd and 3rd lamps were 337, 495 and 756 $\mu\text{mol}/\text{m}^2\text{s}$, respectively.

The CO₂ flow (N), absorbed or rejected, was calculated according to Garry (1988):

$$N = -Q_e \cdot \Delta c [\text{mL}/\text{min}] \quad (2)$$

Where:

Q_e = the airflow passed in the enclosed,

Δc = the difference in CO₂ concentration between the entry and the exit.

The CO₂ flow allowed establishing photosynthesis and respiration regression equations.

Net photosynthesis, P_n , was calculated as follows (Papazoglou *et al.*, 2005):

$$P_n = P \cdot N \cdot \frac{1}{0.0224 \cdot 60 \cdot S} [\mu\text{mol}/\text{m}^2\text{s}] \quad (3)$$

Where:

P = the slope of the regression equation of the function

$\Delta c = f(t)$,

S = the total frond area,

N = the CO₂ flow ($1 \mu\text{mole CO}_2 = 22.4 \times 10 \text{ mL CO}_2$).

Brut photosynthesis (P_b) was expressed as follows:

$$\begin{aligned} P_{b_1} (\text{1st lamp}) &= (P_1 + P_2) \cdot N \\ P_{b_2} (\text{2nd lamp}) &= (P_m + P_2) \cdot N \\ P_b (\text{3rd lamp}) &= (P_2 + P_3) \cdot N \end{aligned} \quad (4)$$

Where:

P_1 and P_2 are the slopes of the regression equation of the 1st and 2nd respiration,

$$m = \frac{p_1 + p_2}{2} \quad (5)$$

P_3 , P_4 and P_5 are the slopes of the regression equation of the 1st, 2nd and 3rd photosynthesis. Net and brut photosynthesis are expressed in $\mu\text{mol}/\text{m}^2\text{s}$.

III. Results

Growth response to copper exposure

Data from experimental dose-response curve (Fig.1) showed that Lemna minor, when exposed to Cu concentrations from 0.3 to 1.0 mg/L, exhibited significant inhibition of growth. For Cu concentration of 0.5 mg/L, duckweed growth was inhibited by 70% (indicated on the figure by the broken line) representing the minimal growth index attained under our experimental conditions.

On the other hand, at concentration ranging from ~ 0 (control) to 0.2 mg/L, the growth index was optimal. Based on these results, Cu concentration of 0.2 mg/L was considered as the threshold toxicity in Lemna minor under conditions indicated above.




Fig. 1: Growth of Lemna minor fronds in the presence of copper in the nutrient medium. Vertical bars indicate standard deviation, n = 3 (Khellaf and Zerdaoui, 2009a)


removal of copper

Fig.2 shows the concentration of copper removed by the plants after 96 h of exposure. The initial Cu concentration in the Coïcand Lesaints solution was 0.2 mg/L. After 1 day, 15% of the initial metal concentration was removed and after 4 days, Lemna minor had removed 26% of Cu from the nutrient medium. The control treatment showed that the aquatic plants were responsible for the disappearance of amount of Cu from water.

photosynthesis and respiration

Effect of copper at 0.2 mg/L on photosynthesis and respiration of Lemna minor is shown in Fig.

3. Dark respiration was inhibited in the presence of Cu; the 1st and 2nd respiration slopes were respectively 0.41 and 0.20, whereas those of the control were 1.70 and 1.66 (Table 1). However, CO_2 assimilation increased in the presence of 0.2 mg/L of cupric ions. Net photosynthesis ranged between 8.3 and 12.2 $\mu\text{mol}/\text{m}^2\text{s}$ for different



Elimination (%)

Fig. 2: Elimination of copper from the nutrient medium in the presence of *L. minor*.

The initial Cu concentration was 0.2 mg/L luminous energies used in this study. Compared to control, net photosynthesis was approximately 3 times higher. Brut photosynthesis was only 2 times of the control because of the inhibition of the gas exchanges in dark. However, for an irradiation $\geq 495 \mu\text{mol/m}^2\text{s}$, it was noticed that saturation was mainly observed in the case of brut photosynthesis (Fig.4).

IV. Discussion

Copper is considered to be one of the most toxic trace metal to plants, although it is required as an essential element for metabolic and physiological processes (Xia and Tian, 2009). Dewezetal. (2005) explained some hypothesis concerning the mechanism of Cu toxicity on the plant growth. Copper was recognized to be a strong inhibitor of photosystem II (PSII) electron transport activity associated to the water splitting system; by this effect, the metal may alter the energy storage via photosynthesis which causes the decrease of biomass growth. However, some plant species tolerate this element at concentrations higher than those used in medium cultures. Our study indicated that, *Lemna minor* was sensitive to copper for concentrations $\geq 0.3 \text{ mg/L}$ and the threshold toxicity was 0.2 mg Cu/L. Published quantitative data for the threshold toxicity of




Fig. 3: CO₂ concentration as a function of time; (a) plants 96 h after exposure to 0.2 mg Cu/L and (b) control values are means of 2 essays

metal ions on duckweed species are very variable (Table 2). This is caused by the different duckweed species used and by the different test conditions, especially concerning the nutrient media as well as by the methods of evaluation (Appenroth et al., 2010).

The Cu removal treatments showed that the aquatic plants were responsible for the disappearance of Cu from water. From earlier results, it seems that metal removal from the medium was due to an accumulation in plants; several studies demonstrated that duckweed species (particularly

Table 1: Slope values of photosynthesis and respiration curves

Gas exchange	Slope values (mg/L. min)			
	Control	R ²	Plants exposed	R ²
Respiration 1	1.70	0.99	0.41	0.80
Respiration 2	1.66	0.97	0.20	0.84
Photosynthesis 1	- 1.50	0.99	- 1.73	0.99
Photosynthesis 2	- 2.18	0.99	- 2.54	0.99
Photosynthesis 3	- 1.83	0.98	- 2.53	0.99

The plants were exposed to 0.2 mg/L during 4 days. 1, 2 and 3 correspond to the 1st, 2nd and 3rd lamp in photosynthesis essays. Different slope values are means of 2 values; error was < 10%. R² is the coefficient of determination.

Lemna minor and Lemnagibba) were able to accumulate elevated amount of Cu in their tissues (Jain et al., 1989; Zayed et al., 1998; Ateret al., 2006; Megatliet al., 2009) inducing an abatement of Cu concentration in water. Our result confirmed that Lemna minor showed a potential of phytoremediation of contaminated waters charged with low concentrations of Cu.

Photosynthetic activities increased in the presence of 0.2 mg/L of cupric ions for different luminous energies. However, according to the study of Olette et al. (2008), copper used as CuSO₄ (pesticide) inhibited the photosynthetic activities of Lemna minor at concentrations of 12, 24, 40 and 100 µg /L. For an exposure time of 7 days, the metal element present in the medium inhibited the photosynthesis of the aquatic plants by 0.4%; this inhibition reached 8% in the presence of 100

µg/L of Cu. Prasad et al., (2001) demonstrated that 1 and 10 µM of Cu present in a Hoagland solution increased (160 and 120% of the control) the concentration of photosynthetic pigments of Lemnatisulca (another species of duckweed). These results agree with those of the present study. The presence of 0.2 mg/L of Cu in the nutrient medium increased considerably the absorption of

Table 2: Threshold toxicity* of copper in duckweed

Duckweed species	Experimental conditions	Threshold toxicity	Reference
Lemna minor	Tap water, pH=7.2	1 mg/L	Jain et al., 1989
Lemna minor	1/4 Hoagland Solution; pH=6	< 5 mg/L	Zayed et al., 1998
Lemna minor	Inorganic growth Medium; pH=6.5	< 0.25 µM	Teisseire and Guy, 2000
Lemna minor	White nutritive Solution; pH=6.8	< 0.5 mg/L	Ateret al., 2006
Lemnagibba	White nutritive Solution; pH=6.8	0.5 mg/L	Ateret al., 2006
Lemnagibba	Hoagland medium; pH=6.5	10 ⁻⁴ mg/L	Megatliet al., 2009
Lemna minor	1/10 Hoagland Solution; pH=6	1.6 mg/L	Kanoun-Bouléet al., 2009

*Threshold concentration is Cu concentration at which no growth inhibition is observed in duckweed biomass.




Fig. 4: Variation of (a) net photosynthesis and (b) brut photosynthesis, with the luminous energy. Treatment corresponds to plants exposed to 0.2 mg/L of Cu. Net and brut photosynthesis were calculated using the values on Table 1.

CO_2 by duckweed. Net photosynthesis increased approximately threetimescomparedtothecontrol. Copper was responsible for 130-290% increases in the photosynthetic activities. This effect might be explained by an increase in electron transport in photosynthetic systems. Copper is an essential elementincellularmetabolism andacatalytic component of proteins and enzymes (Teisseire and Guy, 2000). It is plausible that this element was responsible for the synthesis of plastocyanin (Bertrand and Poirier, 2005). Additionally, this study demonstrated that the increaseinphotosyntheticactivitieswasobserved for different luminous energies. However, foran

irradiation $\geq 495 \mu\text{mo}/\text{m}^2\text{s}$, saturation was mainly noticed in the case of brut photosynthesis (Fig. 4). Beyond that, the capacity of absorption of photons exceeds the capacity of their use. The reactions of CO_2 assimilation become limiting and photosynthetic activities present a maximal intensity.

According to Wedge and Burris (1982) the energy saturation for Lemna minor ranged in the interval $300\text{--}600 \mu\text{mo}/\text{m}^2\text{s}$. Our results showed that the energy saturation for the species used in this study corresponded to a photosynthetic active irradiation of $495\text{--}750 \mu\text{mo}/\text{m}^2\text{s}$.

Finally, it can be concluded that among the tools used to study effects of toxic elements on plants, growth and photosynthesis are often proposed as simple, rapid and sensitive methods. Based on these methods, the results of our study showed that Lemna minor was sensitive to copper for concentrations $\geq 0.3 \text{ mg/L}$. For lowest concentrations, plant growth was optimal and photosynthetic activities increased by 290% under elevated luminous energy. The duckweed species could survive in contaminated medium ($\leq 0.2 \text{ mg/L}$) and could detect sensitively metal concentration $\geq 0.3 \text{ mg/L}$. It was concluded that Lemna minor could be a good candidate for the evaluation of metal pollution in biomonitoring programs for risk assessments and toxic effect prediction.

V. Acknowledgements

The authors thank Prof. J. C. Leclerc (LEPA, Saint- Etienne, France) and Ms. S. Sey (University of Gafsa, Tunisia) for their kind assistance.

References

- [1] Appenroth, K.-J., Krech, K., Keresztes, A., Fischer, W. H., Koloczek, H., (2010). Effects of nickel on the chloroplasts of the duckweeds *Spirodela polyrrhiza* and *Lemna minor* and their possible use in biomonitoring and phytoremediation. *Chemosphere*, **78**:216-223.
- [2] Ater, M., Ait Ali, N., Kasmir, H., (2006). Tolérance et accumulation induite de cuivre et de chrome chez deux espèces de lentilles d'eau: *Lemna minor* L. et *Lemna gibba* L. *Revue des Sciences de l'Eau*, **19** (1) :57-67.
- [3] Bertrand, M., Poirier, I., (2005). Photosynthetic organisms and excess of metals. *Photosynthetica*, **43** (3): 345-353.
- [4] Dewez, D., Geoffroy, L., Vernet, G., Popovic, R., (2005). Determination of photosynthetic and enzymatic Biomarkers sensitivity used to evaluate toxic effects of copper and fludixonil in alga *Scenedesmus obliquus*. *Aquat. Toxicol.*, **74**: 150-159.
- [5] Gary, C., (1988). Prise en compte des différentes sources d'erreurs et estimation de la précision dans la mesure des échanges de CO_2 en système ouvert. *Photosynthetica*, **22** (1): 58-69.
- [6] Hattab, S., Dridi, B., Chouba, L., Ben Kheder, M., Bousetta, H., (2009). Photosynthesis and growth responses of pea *Pisum sativum* L. under heavy metals stress. *J. Environ. Sci.*, **21**:1552-1556.
- [7] Jain, S.K., Vasudevan, P., Jha, N.K., (1989). Removal of some heavy metals from polluted water by aquatic plants: Studies on duckweed and water velvet. *Biol. Wastes*, **28**: 115-126.
- [8] Kanoun-Boulé, M., Vicentea, J.A.F., Nabaisa, C., Prasad, M.N.V., Freitas, F., (2009). Ecophysiological tolerance of duckweeds exposed to copper. *Aquatic Toxicol.*, **91** (1): 1-9.
- [9] Khellaf, N., and Zerdaoui, M., (2009a). Growth response of the duckweed *L. minor* to heavy metals pollution. *Iran. J. Environ. Health Sci. Eng.*, **6** (3): 161-166.
- [10] Khellaf, N., and Zerdaoui, M., (2009b). Phytoaccumulation of zinc by the aquatic plant *Lemna gibba* L. *Bioresour. Technol.*, **100**: 6137-6140.
- [11] Megateli, S., Semsari, S., Couderchet, M., (2009). Toxicity and removal of heavy metals (cadmium, copper and zinc) by *Lemna gibba*. *Ecotoxicol. Environ. Saf.* **72**:1774-1780. Miretzky, P., Saralegui, A., Cirelli, A.F., (2004). Aquatic macrophytes potential for the simultaneous removal of heavy metals (Buenos Aires, Argentina). *Chemosphere*, **57**: 997-1005.
- [12] Movahedian, H., Bina, B., Asghari, G.H., (2005). Toxicity Evaluation of Waste water Treatment Plant Effluents Using *Daphnia magna*. *Iran. J. Environ. Health Sci. Eng.*, **2** (2): 1-4.
- [13] OECD, (2002). Guidelines for the testing of chemicals, Lemna sp. Growth Inhibition Test, Draft guideline 221, Paris, France
- [14] Olette, R., Couderchet, M., Biagiatti, S., Eullaffroy, P., (2008). Toxicity and removal of pesticides by selected aquatic plants. *Chemosphere*, **70** : 1414-1421.
- [15] Papazoglou, E.G., Karantounias, G.A., Vemmos, S.N., Bouranis, D.L., (2005). Photosynthesis and growth responses of Giant reed (*Arundo donax* L.) to heavy metals Cd and Ni. *Environ. Int.*, **31**:243-249.
- [16] Prasad, M.N.V., Malek, P., Waloszek, A., Bojko, M., Strazalka, K., (2001). Physiological responses of *Lemna trisulca* L. (duckweed) to cadmium and copper accumulation. *Plant Sci.*, **161**:881-889.
- [17] Sabreen, S., and Sugiyama, S., (2008). Trade-off between cadmium tolerance and relative growth rate in 10 grass species. *Environ. Experimental Bot.*, **63**: 327-332.
- [18] Teisseire, H., and Guy, V., (2000). Copper induced changes in antioxidant enzymes activities in fronds of duckweed (*Lemna minor*). *Plant Sci.*, **153**: 65-72.
- [19] Vinodhini, R., and Narayanan, M., (2009). The impact of toxic heavy metals on the haematological parameters in common carp (*Cyprinus carpio* L.). *Iran. J. Environ. Health Sci. Eng.*, **6** (1):23-28.
- [20] Wedge, R.M., and Burris, J.E., (1982). Effects of light and temperature on duckweed photosynthesis. *Aquat. Bot.*, **13**: 133-140.
- [21] Xia, J., and Tian, Q., (2009). Early stage toxicity of excess copper to photosystem II of *Chlorella pyrenoidosa*—OJIP chlorophyll a fluorescence analysis. *J. Environ. Sci.*, **21**:1569-1574.
- [22] Zayed, A., Gowthaman, S., Terry, N., (1998). Phytoaccumulation of trace elements by wetland plants: I- Duckweed. *J. Environ. Qual.*, **27**: 715-721.
- [23] Zhou, Q., Zhang, J., Fu, J., Shi, J., Jiang, G., (2008). Biomonitoring: An appealing tool for assessment of metal pollution in the aquatic ecosystem. *Analytica Chimica Acta*, **606**: 135-150.